

NOAA Technical Report NMFS Circular 413

Marine Flora and Fauna of the Northeastern United States. Crustacea: Branchiura



Roger F. Cressey

May 1978



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Circulars

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS Circular series continues a series that has been in existence since 1941. The Circulars are technical publications of general interest intended to aid conservation and management. Publications that review in considerable detail and at a high technical level certain broad areas of research appear in this series. Technical papers originating in economics studies and from management investigations appear in the Circular series.

NOAA Technical Report NMFS Circulars are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D825, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent Circulars are:

365. Processing EASTROPAC STD data and the construction of vertical temperature and salinity sections by computer. By Forrest R. Miller and Kenneth A. Bliss. February 1972, iv + 17 p., 8 figs., 3 app. figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

366. Key to field identification of anadromous juvenile salmonids in the Pacific Northwest. By Robert J. MacConnell and George R. Snyder. January 1972, iv + 6 p., 4 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

367. Engineering economic model for fish protein concentration processes. By K. K. Almenas, L. C. Durilla, R. C. Ernst, J. W. Gentry, M. B. Hale, and J. M. Marchello. October 1972, iii + 175 p., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

368. Cooperative Gulf of Mexico estuarine inventory and study, Florida: Phase I, area description. By J. Kneeland McNulty, William N. Lindall, Jr., and James E. Sykes. November 1972, vii + 126 p., 46 figs., 62 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

369. Field guide to the angelfishes (Pomacanthidae) in the western Atlantic. By Henry A. Feddern. November 1972, iii + 10 p., 17 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

370. Collecting and processing data on fish eggs and larvae in the California Current region. By David Kramer, Mary J. Kalin, Elizabeth G. Stevens, James R. Thrailkill, and James R. Zweifel. November 1972, iv + 38 p., 38 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

371. Ocean fishery management: Discussion and research. By Adam A. Sokoloski (editor). (17 papers, 24 authors.) April 1973, vi + 173 p., 38 figs., 32 tables, 7 app. tables.

372. Fishery publications, calendar year 1971: Lists and indexes. By Thomas A. Manar. October 1972, iv + 24 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

374. Marine flora and fauna of the northeastern United States. Annelida: Oligochaeta. By David G. Cook and Ralph O. Brinkhurst. May 1973, iii + 23 p., 82 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

375. New Polychaeta from Beaufort, with a key to all species recorded from North Carolina. By John H. Day. July 1973, xiii + 140 p., 18 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

376. Bottom-water temperatures on the continental shelf, Nova Scotia to New Jersey. By John B. Colton, Jr. and Ruth R. Stoddard. June 1973, iii + 55 p., 15 figs., 12 app. tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

377. Fishery publications, calendar year 1970: Lists and indexes. By Mary Ellen Enggett and Lee C. Thorson. December 1972, iv + 34 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

378. Marine flora and fauna of the northeastern United States. Protozoa: Ciliophora. By Arthur C. Borror. September 1973, iii + 62 p., 5 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

379. Fishery publications, calendar year 1969: Lists and indexes. By Lee C. Thorson and Mary Ellen Enggett. April 1973, iv + 31 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

380. Fishery publications, calendar year 1968: Lists and indexes. By Mary Ellen Enggett and Lee C. Thorson. May 1973, iv + 24 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

381. Fishery publications, calendar year 1967: Lists and indexes. By Lee C. Thorson and Mary Ellen Enggett. July 1973, iv + 22 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

382. Fishery publications, calendar year 1966: Lists and indexes. By Mary Ellen Enggett and Lee C. Thorson. July 1973, iv + 19 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

383. Fishery publications, calendar year 1965: Lists and indexes. By Lee C. Thorson and Mary Ellen Enggett. July 1973, iv + 12 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

384. Marine flora and fauna of the northeastern United States. Higher plants of the marine fringe. By Edwin T. Moul. September 1973, iii + 60 p., 109 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

385. Fishery publications, calendar year 1972: Lists and indexes. By Lee C. Thorson and Mary Ellen Enggett. November 1973, iv + 23 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

386. Marine flora and fauna of the northeastern United States. Pycnogonida. By Lawrence R. McCloskey. September 1973, iii + 12 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

387. Marine flora and fauna of the northeastern United States. Crustacea: Stomatopoda. By Raymond B. Manning. February 1974, iii + 6 p., 10 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.



NOAA Technical Report Circular 413

**Marine Flora and Fauna of
the Northeastern United States.
Crustacea: Branchiura**

Roger F. Cressey

May 1978

U.S. DEPARTMENT OF COMMERCE

Juanita M. Kreps, Secretary

National Oceanic and Atmospheric Administration

Richard A. Frank, Administrator

National Marine Fisheries Service

FOREWORD

This issue of the "Circulars" is part of a subseries entitled "Marine Flora and Fauna of the Northeastern United States." This subseries will consist of original, illustrated, modern manuals on the identification, classification, and general biology of the estuarine and coastal marine plants and animals of the northeastern United States. Manuals will be published at irregular intervals on as many taxa of the region as there are specialists available to collaborate in their preparation.

The manuals are an outgrowth of the widely used "Keys to Marine Invertebrates of the Woods Hole Region," edited by R. I. Smith, published in 1964, and produced under the auspices of the Systematics-Ecology Program, Marine Biological Laboratory, Woods Hole, Mass. Instead of revising the "Woods Hole Keys," the staff of the Systematics-Ecology Program decided to expand the geographic coverage and bathymetric range and produce the keys in an entirely new set of expanded publications.

The "Marine Flora and Fauna of the Northeastern United States" is being prepared in collaboration with systematic specialists in the United States and abroad. Each manual will be based primarily on recent and ongoing revisionary systematic research and a fresh examination of the plants and animals. Each major taxon, treated in a separate manual, will include an introduction, illustrated glossary, uniform originally illustrated keys, annotated check list with information when available on distribution, habitat, life history, and related biology, references to the major literature of the group, and a systematic index.

These manuals are intended for use by biology students, biologists, biological oceanographers, informed laymen, and others wishing to identify coastal organisms for this region. In many instances the manuals will serve as a guide to additional information about the species or the group.

Geographic coverage of the "Marine Flora and Fauna of the Northeastern United States" is planned to include organisms from the headwaters of estuaries seaward to approximately the 200-m depth on the continental shelf from Maine to Virginia, but may vary somewhat with each major taxon and the interests of collaborators. Whenever possible representative specimens dealt with in the manuals will be deposited in the reference collections of major museums in the region.

After a sufficient number of manuals of related taxonomic groups have been published, the manuals will be revised, grouped, and issued as special volumes. These volumes will thus consist of compilations of individual manuals within phyla such as the Cnidaria, Arthropoda, and Mollusca, or of groups of phyla.

CONTENTS

Introduction	1
Glossary	1
Key to the Branchiura of northeastern United States	3
Annotated systematic list	8
Selected bibliography	8
Systematic index	9
Acknowledgments	10
Coordinator's comments	10

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

Marine Flora and Fauna of the Northeastern United States. Crustacea: Branchiura

ROGER F. CRESSEY¹

ABSTRACT

Eleven species of *Argulus* are known from the northeastern United States. An illustrated key and an annotated list of these species with notes on their hosts and distribution within and without the study area are included. New host records are included.

INTRODUCTION

The class Branchiura, once considered a subgroup of the class Copepoda, consists of a single order (Argulidea) and two families (Argulidae and Dipteropeltidae); the first with three genera (*Argulus*, *Chonopeltis*, and *Dolops*) and the latter with two (*Dipteropeltis* and *Talaus*). *Argulus* is cosmopolitan in distribution and the only genus reported from North America and Europe.

Members of the class (commonly known as "fish lice") are parasitic on freshwater and coastal marine fishes. They are characterized by the presence of compound eyes, prehensile antennae, and the second maxillae transformed into sucker disks (except *Dolops*). The feeding mechanism consists of a long preoral stylet or "string" and a posterior mouth tube (Fig. 1). The parasite repeatedly pierces the host tissue with the stylet and probably uses the mouth tube to ingest fluids oozing from the resultant wounds.

Eggs are attached to substrate objects (rocks, vegetation, etc.). Development is direct (newly hatched forms resemble the adult). Tokiota (1936) described seven larval stages and several subsequent molts for *Argulus japonicus*, and he reported that sexual maturity was achieved in about 30 days. The sexes can be distinguished in the first larval stage. Adult males are generally smaller than females and can be distinguished from them by the presence of a pair of elongate testes within the abdomen along each side of the midline and by the presence of sclerotized processes on the basal segments of thoracic legs 2, 3, and 4. Females possess a pair of rounded spermathecae in the proximal third of the abdomen; the basal portions of the legs are unmodified.

Argulids are "loosely" associated with their hosts and quickly drop off when the fish is removed from water. Consequently, fish should be examined as soon after capture as possible. The parasites are found on the body surface, in the gill chamber, and in the mouth of the host. Argulids are often collected in plankton samples as well, due to their transient attachment to their hosts. Collections should be preserved in 70% ethyl alcohol; 5% Formalin is acceptable for short-term preservation. Where Formalin is used, the material should be transferred to alcohol as soon as possible as Formalin tends to decalcify materials over a long period of time. It may be necessary to clear heavily pigmented specimens in a warm solution of 10% KOH for 15 min or more in order to study them microscopically. Small or less pigmented specimens may be cleared in lactic acid for study. To examine the respiratory area it may be necessary to dissect one side of the carapace from the rest of the body.

GLOSSARY

- Carapace* dorsolateral plate covering cephalothorax.
Respiratory areas two areas on each ventrolateral part of the carapace devoid of sclerotization and open to the overlying tissues.
Scales short integumental processes with blunt or broadly rounded tips occurring singly or in groups.
Spines short integumental processes with pointed tips occurring singly or in groups.
Sucker modified second maxilla, seen ventrally, one on each side of the oral area.
Sucker rod supporting structures composed of a series of sclerotized processes radiating outwardly in the membrane fringing the outer edge of the sucker.

¹Smithsonian Institution, U.S. National Museum of Natural History, Washington, DC 20560.




Figure 1.—*Argulus alosae*, female, ventral view, labeled.

KEY TO THE BRANCHIURA OF NORTHEASTERN UNITED STATES

- 1 Smaller respiratory area mostly or entirely anterior to larger (specimens "mostly anterior" have sucker rods of 5-7 elements)2




Figure 2.—Respiratory areas of: a. *Argulus japonicus*; b. *A. chesapeakensis*; c. *A. laticauda*; d. *A. megalops*; e. *A. alosae*; f. *A. funduli*.

- 1 Smaller respiratory area mostly or entirely lateral to larger (specimens "mostly lateral" have sucker rods of 2 elements)7




Figure 3.—Respiratory areas of: a. *Argulus versicolor*; b. *A. catastomi*; c. *A. appendiculosus*; d. *A. stizostethi*; e. *A. maculosus*.

2 (1)	Mouth tube with spines or scales at base3
2 (1)	Mouth tube without spines or scales at base5




Figure 4.—Mouth tubes of: a. *Argulus japonicus*; b. *A. alosae*; c. *A. megalops*.

3 (2)	More than 15 elements in sucker rods	<i>Argulus funduli</i>
-------	--	------------------------




Figure 5.—Sucker rods of *Argulus funduli*.

3 (2)	Less than 15 elements in sucker rods4
-------	--	----

4 (3)	Mouth tube with spines at base (see Fig. 4b). Sucker rods as in Figure 6	<i>Argulus alosae</i>
-------	--	-----------------------





Figure 6.—Sucker rods of *Argulus alosae*.

Figure 7.—Sucker rods of *Argulus megalops*.



- 4 (3) Mouth tube with scales at base (see Fig. 4c).
Sucker rods as Figure 7 *Argulus megalops*

- 5 (2) More than 5 elements in sucker rods . . . *Argulus japonicus*





Figure 8.—Sucker rods of *Argulus japonicus*.

- 5 (2) Less than 6 elements in sucker rods 6


- 6 (5) Antennal spines nearly as wide as long. Sucker rods of 4-5 elements *Argulus laticauda*

Figure 9.—*Argulus laticauda*: a. antennae; b. sucker rods.



- 6 (5) Antennal spines much longer than wide. Sucker rods of 3-4 elements *Argulus chesapeakeensis*


Figure 10.—*Argulus chesapeakeensis*: a. antennae; b. sucker rods.



7 (1)	More than 8 elements in sucker rods8
7 (1)	Less than 8 elements in sucker rods9


8 (7)	More than 12 elements in sucker rods	<i>Arbulus stizostethi</i>
-------	--	----------------------------

Figure 11.—Sucker rods of *Argulus stizostethi*.



8 (7)	Less than 12 elements in sucker rods	<i>Argulus catostomi</i>
-------	--	--------------------------

Figure 12.—Sucker rods of *Argulus catostomi*.



- 9 (7) More than 3 elements in sucker rods
 *Argulus versicolor*




Figure 13.—Sucker rods of *Argulus versicolor*.

- 9 (7) Less than 3 elements in sucker rods 10

- 10 (9) Basal element of sucker rod longer than terminal
 *Argulus maculosus*




Figure 14.—Sucker rods of *Argulus maculosus*.

- 10 (9) Basal element of sucker rod
 shorter than terminal . . .
 *Argulus appendiculosus*




Figure 15.—Sucker rods of *Argulus appendiculosus*.

ANNOTATED SYSTEMATIC LIST

The 11 species of *Argulus* known from the north-eastern United States are arranged alphabetically. Notes on the distribution and hosts of each species within and without the study area are included. Comments on ecology and frequency are included wherever appropriate. An asterisk denotes new host record.

Argulus alosae Gould 1841. Distribution: east coast of North America from Nova Scotia to the Gulf coast of Texas. Hosts: *Microgadus tomcod*; *Tautogolabrus adspersus*; *Alosa pseudoharengus*; *Dorosoma cepedianum*; *Clupea harengus*; *Cynoscion nebulosus*; *Gasterosteus* sp. [originally reported as *G. bispinosus* from the Gulf of St. Lawrence (Wilson, 1902)]; *Strömgyllura marina*; *Rhopilema verrilli** (medusa). This species has been reported from several species of inshore marine fish but its repeated occurrence on clupeids (*Alosa*, *Clupea*, and *Dorosoma*) may indicate a preference for that host family.

Argulus appendiculosus Wilson 1907. Distribution: northern half of United States from Vermont to Virginia west to Wyoming and also Texas and Louisiana. Hosts: *Catostomus catostomus*; *Ictiobus cyprinellus*, *I. bubalus*; *Ictalurus punctatus*, *I. melas*, *I. nebulosus*; *Amia calva*; *Micropterus salmoides*; *Morone (Roccus) chrysops*; *Perca flavescens*; *Cyprinus carpio*; *Stizostedion vitreum*; *Dorosoma cepedianum*; *Lepisosteus osseus**. *Argulus appendiculosus* has so far only been reported from freshwater, but since one of its hosts (*Dorosoma*) is found in coastal marine waters as well, the parasite may also be eventually found there.

Argulus catostomi Dana and Herrick 1837. Distribution: northern United States from Massachusetts west to Minnesota. Hosts: *Catostomus commersoni*, *C. catostomus*; *Hypentelium nigricans*; *Erimyzon sucetta*; *Cyprinus carpio*; *Notemigonus crysoleucas*. So far this parasite has been found only in freshwater and on members of the families Catostomidae and Cyprinidae.

Argulus chesapeakeensis Cressey 1971. Distribution: southeast coast of United States from Chesapeake Bay south to Sapelo Island, Ga. Hosts: *Opsanus tau*; *Archosargus probatocephalus**; *Arius felis**; *Mugil cephalus**; *Gobiosoma boscii**; *Paralichthys dentatus*; *Dasyatis americana**; *Pteroplatea maclura**; *Rachycentron canadum*. This species has so far been collected only from the southeast coast of the United States. It is closely related to *A. laticauda*, found along the northeast coast.

Argulus funduli Kroyer 1863. Distribution: east coast of North America from New Brunswick, Canada, south to the mouth of the Neches River, Texas, collected at several localities between. Hosts: *Menidia notata*; *Lagodon rhomboides*; *Pseudopleuronectes americanus*; *Cyprinodon variegatus*; *Fundulus grandis*, *F. heteroclitus*, *F. majalis*, *F. ocellaris*. This species is common along the entire east coast of the United

States and is most often associated with marine cyprinodontids.

Argulus japonicus Thiele 1900. Distribution: entire United States. Hosts: *Carassius auratus* (goldfish). This argulid is an introduced species brought into the United States from the Orient on aquarium fish. It is very common anywhere goldfish are found. No records from marine waters to date.

Argulus laticauda Smith 1873. Distribution: northeast coast of United States from New England south to Long Island Sound. Hosts: *Prionotus* sp.; *Anguilla rostrata*; *Pseudopleuronectes americanus*. Common on the American eel and winter flounder of the northeast coast.

Argulus maculosus Wilson 1902. Distribution: eastern half of United States and southeastern Canada as far west as Iowa and collected in northern and southernmost states. Hosts: *Ictalurus natalis*, *I. nebulosus*; *Amia calva*; *Ambloplites rupestris*; *Erimyzon sucetta*; *Esox* sp.; *Umbra limi*; *Lepisosteus osseus**. This common argulid has so far been collected only in freshwater.

Argulus megalops Smith 1873. Distribution: east coast of North America from New Brunswick, Canada, to Florida. Hosts: a wide variety of inshore marine fishes (recorded from 16 species) with no indications as to host preferences.

Argulus stizostethi Kellicott 1880. Distribution: north-eastern quarter of the United States from Iowa and Minnesota east to New England and New Brunswick, Canada. Hosts: *Salvelinus fontinalis*; *Stizostedion vitreum*, *S. canadense*; *Acipenser fulvescens*; *Esox masquinongy*; *Alosa sapidissima*; *Gasterosteus* sp.; *Notropis* sp.; *Coregonus* sp.; *Dorosoma cepedianum**. This parasite seems to be restricted to those states near the Great Lakes and St. Lawrence River. All collections are from freshwater except that from *Dorosoma* from Woods Hole, Mass.

Argulus versicolor Wilson 1902. Distribution: scattered throughout the eastern half of the United States. Recorded in Massachusetts, Maryland, Indiana, Georgia, and Texas. Hosts: *Esox niger*; *Ambloplites* sp.; and "perch." So far there are no marine records for this species, but four of the recorded states are marine coastal.

SELECTED BIBLIOGRAPHY

CRESSEY, R. F.

1972. The genus *Argulus* (Crustacea: Branchiura) of the United States. Bioat of Freshwater Ecosystems. Environmental Protection Agency, Identification Manual No. 2, 14 p.

MEEHAN, O. L.

1940. A review of the parasitic Crustacea of the genus *Argulus* in the collections of the United States National Museum. Proc. U.S. Nat. Mus. 88:459-522.

TOKIOKA, T.

1936. Larval development and metamorphosis of *Argulus japonicus*. Mem. Coll. Sci. Kyoto Imp. Univ. ser. B, 12:93-114.

WILSON, C. B.

1903. North American parasitic copepods of the family Argulidae with a bibliography of the group and a systematic review of all known species. Proc. U.S. Nat. Mus. 25:635-742.

SYSTEMATICS INDEX

<i>Acipenser fulvescens</i>	8	<i>Esox</i>	
<i>Alosa</i>	8	<i>masquinongy</i>	8
<i>pseudoharengus</i>	8	<i>niger</i>	8
<i>sapidissima</i>	8	sp.	8
<i>Ambloplites</i>		<i>Fundulus</i>	
<i>rupestris</i>	8	<i>grandis</i>	8
sp.	8	<i>heteroclitis</i>	8
<i>Amia calva</i>	8	<i>majalis</i>	8
<i>Anguilla rostrata</i>	8	<i>ocellaris</i>	8
<i>Archosargus probatocephalus</i>	8	<i>Gasterosteus</i> sp.	8
<i>Argulidae</i>	1	<i>Gobiosoma bosci</i>	8
<i>Argulus</i>		<i>Hypentelium nigricans</i>	8
<i>allosae</i>	2, 3, 4, 8	<i>Ictalurus</i>	
<i>appendiculosus</i>	3, 7, 8	<i>melas</i>	8
<i>catostomi</i>	3, 6, 8	<i>natalis</i>	8
<i>chesapeakeensis</i>	3, 5, 8	<i>nebulosus</i>	8
<i>funduli</i>	3, 4, 8	<i>punctatus</i>	8
<i>japonicus</i>	1, 3, 4, 5, 8	<i>Ictiobus</i>	
<i>laticauda</i>	3, 5, 8	<i>bubalus</i>	8
<i>maculosus</i>	3, 7, 8	<i>cyprinellus</i>	8
<i>megalops</i>	3, 4, 5, 8	<i>Lagodon rhomboides</i>	8
<i>stizostethi</i>	3, 6, 8	<i>Lepisosteus osseus</i>	8
<i>versicolor</i>	3, 7, 8	<i>Menidia notata</i>	8
<i>Arius felis</i>	8	<i>Microgadus tomcod</i>	8
<i>Carassius auratus</i>	8	<i>Micropterus salmoides</i>	8
<i>Catostomidae</i>	8	<i>Morone chrysops</i>	8
<i>Catostomus</i>		<i>Mugil cephalus</i>	8
<i>catostomus</i>	8	<i>Notemigonus crysoleucas</i>	8
<i>commersoni</i>	8	<i>Notropis</i> sp.	8
<i>Chonopeltis</i>	1	<i>Opsanus tau</i>	8
<i>Clupea</i>	8	<i>Paralichthys dentatus</i>	8
<i>harengus</i>	8	<i>Perca flavescens</i>	8
<i>Coregonus</i> sp.	8	<i>Prionotus</i> sp.	8
<i>Cynoscion nebulosus</i>	8	<i>Pseudopleuronectes americanus</i>	8
<i>Cyprinidae</i>	8	<i>Pteroplatea maclura</i>	8
<i>Cyprinodon variegatus</i>	8	<i>Rachycentron canadum</i>	8
<i>Cyprinus carpio</i>	8	<i>Rhopilema verrilli</i>	8
<i>Dasyatis americana</i>	8	<i>Salvelinus fontinalis</i>	8
<i>Dipteropeltidae</i>	1	<i>Stizostedion</i>	
<i>Dipteropeltis</i>	1	<i>canadense</i>	8
<i>Dolops</i>	1	<i>vitreum</i>	8
<i>Dorosoma</i>	8	<i>Strongylura marina</i>	8
<i>cepedianum</i>	8	<i>Talaus</i>	1
<i>Erimyson sucetta</i>	8	<i>Tautoglabrus adspersus</i>	8
		<i>Umbra limi</i>	8

ACKNOWLEDGMENTS

Preparation of the "Marine Flora and Fauna of the Northeastern United States" is being coordinated by the following Board:

Coordinating Editor:

Melbourne R. Carriker, College of Marine Studies, University of Delaware, Lewes, DE 19958.

Editorial Advisers:

Marie B. Abbott, Marine Biological Laboratory, Woods Hole, Mass.

Arthur G. Humes, Boston University Marine Program, Marine Biological Laboratory, Woods Hole, Mass.

Wesley N. Tiffney, Department of Biology, Boston University, Boston, Mass.

Ruth D. Turner, Museum of Comparative Zoology, Harvard University, Cambridge, Mass.

Roland L. Wigley, National Marine Fisheries Service
Northeast Fisheries Center, NOAA, Woods Hole, Mass.
Robert T. Wilce, Department of Botany, University of Massachusetts, Amherst, Mass.

The Board, which established the format for the "Marine Flora and Fauna of the Northwestern United States," invites systematists to collaborate in the preparation of manuals, reviews manuscripts, and advises the Scientific Editor of the National Marine Fisheries Service.

The illustrations were done by Hillary Boyle. The manuscript was critically reviewed by Roland Wigley and Arthur Humes. The materials and records embodied in this paper are housed in the National Museum of Natural History, Washington, D.C.

COORDINATING EDITOR'S COMMENTS

Publication of the "Marine Flora and Fauna of the Northeastern United States" is most timely in view of the growing universal emphasis on environmental work and the urgent need for more precise and complete identification of coastal organisms than has been available. It is mandatory, whenever possible, that organisms be identified accurately to species. Accurate scientific names unlock the great quantities of biological information stored in libraries, obviate duplication of research already done, and often make possible prediction of attributes of organisms that have been inadequately studied.

Roger Cressey started working with parasitic copepods and branchiurans as a graduate student at Boston University in 1958. After receiving his doctoral degree in 1965, he joined the staff of the National Museum of Natural History, Smithsonian

Institution. His studies on parasitic crustaceans have resulted in over 40 publications. He has conducted field work in the Indian Ocean, Madagascar, southeastern Pacific Ocean, and Gulf of Mexico off the west coast of Florida.

Preparation of this manual was supported in part by a grant from the Environmental Protection Agency to the Editorial Board of the "Marine Flora and Fauna of the Northeastern United States." Work on the "Marine Flora and Fauna of the Northeastern United States" by the Coordinating Editor is supported by the College of Marine Studies, University of Delaware.

Manuals are available for purchase from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402. The manuals so far published in the series are listed below:

Marine Flora and Fauna of the Northeastern United States:

Circular

COOK, DAVID G., AND RALPH O. BRINKHURST. Annelida: Oligochaeta.	374
BORROR, ARTHUR C. Protozoa: Ciliophora.	378
MOUL, EDWIN T. Higher Plants of the Marine Fringe.	384
McCLOSKEY, LAWRENCE R. Pycnogonida.	386
MANNING, RAYMOND B. Crustacea: Stomatopoda.	387
WILLIAMS, AUSTIN B. Crustacea: Decapoda.	389
POLLOCK, LELAND W. Tardigrada.	394
LARSON, RONALD J. Cnidaria: Scyphozoa.	397
CAVALIERE, A. R. Higher Fungi: Ascomycetes, Deuteromycetes, and Basidiomycetes.	398
COULL, BRUCE C. Copepoda: Harpacticoida.	399
CUTLER, EDWARD B. Sipuncula.	403
PAWSON, DAVID L. Echinodermata: Holothuroidea.	405
HO, JU-SHEY. Copepoda: Lernaepodida.	406
HO, JU-SHEY. Copepoda: Cyclopoids Parasitic on Fishes.	
CRESSEY, ROGER F. Crustacea: Branchiura.	

388. Proceedings of the first U.S.-Japan meeting on aquaculture at Tokyo, Japan, October 18-19, 1971. William N. Shaw (editor). (18 papers, 14 authors.) February 1974, iii + 133 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

389. Marine flora and fauna of the northeastern United States. Crustacea: Decapoda. By Austin B. Williams. April 1974, iii + 50 p., 111 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

390. Fishery publications, calendar year 1973: Lists and indexes. By Mary Ellen Engett and Lee C. Thorson. September 1974, iv + 14 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

391. Calanoid copepods of the genera *Spinocalanus* and *Mimocalanus* from the central Arctic Ocean, with a review of the Spinocalanidae. By David M. Damkaer. June 1975, x + 88 p., 225 figs., 4 tables. For sale

by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

392. Fishery publications, calendar year 1974: Lists and indexes. By Lee C. Thorson and Mary Ellen Engett. June 1975, iv + 27 p., 1 fig.

393. Cooperative Gulf of Mexico estuarine inventory and study—Texas: Area description. By Richard A. Diener. September 1975, vi + 129 p., 55 figs., 26 tables.

394. Marine Flora and Fauna of the Northeastern United States. Tardigrada. By Leland W. Pollock. May 1976, iii + 25 p., figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

395. Report of a colloquium on larval fish mortality studies and their relation to fishery research, January 1975. By John R. Hunter. May 1976, iii + 5 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.



A000072018910

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

NOAA, the *National Oceanic and Atmospheric Administration*, was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth, and to assess the socioeconomic impact of natural and technological changes in the environment.

The six Major Line Components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

TECHNICAL SERVICE PUBLICATIONS—These are publications containing data, observations, instructions, etc. A partial listing: Data serials; Prediction and outlook periodicals; Technical manuals, training papers, planning reports, and information serials; and Miscellaneous technical publications.

ATLAS—Analysed data generally presented in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER
ENVIRONMENTAL DATA SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

**3300 Whitehaven Street, N.W.
Washington, D.C. 20235**